CLAIMS

- 1. A method for preparing a wine, characterized in that yeast previously enriched in glutathione is introduced into the must at the beginning of, during or after the alcoholic fermentation step.
- 2. The method as claimed in claim 1, characterized in that said glutathione-enriched yeast contains more than 0.5% of glutathione by weight relative to the weight of solids of the yeast.
- 3. The method as claimed in claim 1, characterized in that said glutathione-enriched yeast contains at least 1% of glutathione, and preferably at least 1.5% of glutathione, by weight relative to the weight of solids of the yeast.
- 4. The method as claimed in one of claims 1 to 3, characterized in that said glutathione-enriched yeast is introduced into the must in a proportion of 0.1 g to 1 g of solids per liter of must, preferably in a proportion of 0.3 g/l to 0.7 g/l, and more preferably in a proportion of 0.5 g/l.

25

30

10

- 5. The method as claimed in one of the preceding claims, characterized in that the introduction of said glutathione-enriched yeast into the must produces a provision of at least 3 mg of glutathione per liter of must, preferably at least 5 mg, and more preferably approximately 9 mg of glutathione per liter of must.
- 6. The method as claimed in any one of the preceding claims, characterized in that said glutathione-enriched yeast is introduced into the must in one or more of the following forms: active dry yeast, inactive dry yeast.

- 7. The method as claimed in one of the preceding claims, characterized in that said glutathione-enriched yeast consists at least in part of the yeast inoculated into the must in order to carry out the alcoholic fermentation.
- 8. A must undergoing fermentation or after fermentation, characterized in that it comprises glutathione-enriched yeasts.

10

5

- 9. The must as claimed in the preceding claim, characterized in that it comprises yeasts enriched at more than 0.5% in glutathione, preferably at least 1%, and more preferably at least 1.5%, by weight of
- 15 glutathione relative to the weight of solids of the yeast.
- 10. The must as claimed in either of claims 8 or 9, characterized in that it comprises an amount of glutathione-enriched yeasts corresponding to a content of at least 3 mg of glutathione per liter of must, preferably at least 5 mg, and more preferably approximately 9 mg of glutathione per liter of must.
- 25 11. The use of the method as claimed in one of claims 1 to 7 for the preparation of a wine, in particular of a white wine.
- 12. The use of the method as claimed in one of claims
 30 1 to 7 for the prevention of the browning of white wines after bottling.
 - 13. A wine prepared by means of a method as claimed in one of claims 1 to 7.

35

14. A wine produced from a must as claimed in one of claims 8 to 10.